



Body Condition Scoring Guidance

Corresponds with AW1.3

Introduction

The RMS requires that the body condition of goats is routinely monitored and recorded as part of the farm's management system to confirm health of the goat. This document provides guidance on how to conduct Body Condition Scoring for goats. The assessment guidance is also applicable to RMS auditors.

Please note: Images and links in this guidance refer to sheep. The scoring system is the same for angora goats but due to different fat deposits they will generally score lower. Angora goat specific guidance videos will be added to the guidance shortly.

Description

Body condition scoring is a standardized method to estimate the amount of fat on a goat's body. The body condition score measures the balance between intake and expenditure of energy and is known to be related to feeding motivation. Body condition can be affected by a variety of factors such as food availability, reproductive or productive status, weather conditions, parasites, dental problems, diseases and feeding practices.

How to assess (individual)

Although it may be possible to get some idea of Body Condition by visual assessment of a goat, the most accurate assessment requires goats to be handled. Body Condition Score (BCS) should be assessed in a restrained goat in a race. Body condition is assessed by palpation of the spine in the lumber region just after the last rib. Feel for the horizontal and vertical processes and assess the amount of fat and muscle overlying the bones.








Illustration from Beef and Lamb New Zealand



There are several videos available online demonstrating how to carry out BCS.

How to condition score a ewe (Beef and Lamb New Zealand):

https://www.youtube.com/watch?v=I2 27XYEUOo&index=6&list=PL9ZU9GuQ1pFZhnvT4Wy1aBcxUI4cy 09-n

Body Condition Scoring Demo (Beef and Lamb New Zealand):

https://www.youtube.com/watch?v=CrWQJ7B-ZMQ

How to condition score sheep (DAFWA) (YouTube):


https://www.youtube.com/watch?v=1F5V-GcG1Qk

How to assess (flock)


Randomly draft 25-50 goats into a race or choose a random group from the middle of the flock. A couple of animals from each race when carrying out drenching or other animal husbandry tasks can also be selected. The animal should be standing in a relaxed position and should not be tense or crushed by other animals. Each assessment should only take a matter of seconds.

How to score


The BCS described by Russell et al. (1969, J Agric Sci, 72, 451-454) can be used. For welfare purposes animals are considered thin if they score below 2.0 on this scale, emaciated if they are at or below 1.0, and fat if they are above 4.0. This system is used for all goat breeds and all purposes of use.











Key scores

Optimum BCS will vary across the production cycle and breed of goat used, but in general does that score between 2.5 and 3.5 will provide the optimum production and profitability. Low (<2) and high (>4) BCS have the potential to negatively impact production factors such as ovulation rate, embryonic loss, conception rates and return to service and also carry a higher risk of pregnancy toxemia. Does that are too thin (<2.5) will have poor production, increased risk of mortality and poor reproduction. No goat should score less than 2 without action being taken. The recommended BCS for different production stages are as follows:

- Breeding BCS 2.5 to 3.5
- Early pregnancy maintain BCS at 2.5 to 3.5
- Kidding BCS 2.5 to 3.0, with an absolute minimum of 2.0
- Weaning BCS minimum 2.0.

How often to assess BCS

AW1.3 requires that goats are routinely monitored for BCS. It is recommended that goat have their BCS assessed at least three times during the production cycle, for example as follows:





- When kids are weaned; to allow for does to be managed appropriately for the expected gain in BCS in the run up to breeding.
- In mid-pregnancy, at least for does bearing multiple kids, to allow for any adjustment in feeding for does in poor condition prior to the period of increased nutritional need in late pregnancy.
- Prior to kidding so that does of lower BCS can be offered supplemental feed in lactation if necessary.

How to record BCS

A random sample from the flock can be used to get an average condition score that can be used to help inform decision making. A simple chart (see example below) can be used to record the BCS of a group and any shifts that occurs between recording sessions. Record the body condition of each goat with an X on the chart. The median score of the flock is the score at the middle of the distribution. A printable chart is available in the Records section.

			X			
			X			
		Х	Х			
		Х	Х			
	Х	Х	X	Х		